

croscopy, scanning tunnelling microscopy at ambient conditions, combined with TEM and electron backscatter diffraction. This enables the observation of several important features: Nanostripes are formed by chains of clusters, representing the LRE/Ba substitution. The periodicity of the nanostripes is found to range between 40 and 60 nm;

the shape of the nanoclusters is elliptic with a major axis length between 200 and 500 nm and a minor axis length of about 30 to 150 nm. The dimensions of the nanostripes are similar for both types of NEG samples.

TT 28: Superconductivity: Heterostructures, Andreev Scattering, Proximity Effect, Coexistence

Time: Wednesday 15:15–18:00

Location: HSZ 105

Invited Talk TT 28.1 Wed 15:15 HSZ 105
Unconventional Superconductivity induced by Interfaces and Surfaces — ●MATTHIAS ESCHRIG — Institut für Theoretische Festkörperphysik and DFG-Center for Functional Nanostructures, Universität Karlsruhe, D-76128 Karlsruhe, Germany

Ordered many-body states in solids are often characterized by an order parameter that breaks one or more of the symmetries of the crystal. Such unconventional states lead to interesting new physics associated with the spontaneously broken symmetries. However, in order that such a symmetry breaking can occur it has to be energetically favored. Some of the most interesting symmetry broken states have never been found experimentally in bulk materials for that reason.

However, symmetries can be broken also by introducing interfaces with other materials. In this case, the evasive unconventional states might be induced locally near the interface, and can then penetrate as correlations into bulk materials. The properties of the induced states depend on the scattering characteristics of the interfaces and on the proximity induced states produced by the adjacent materials.

We discuss in particular interface-induced unconventional superconductivity in heterostructures with magnetically active materials, that may exhibit e.g. odd-frequency pairing or equal-spin triplet pairing states. We study the conditions under which such unconventional pairing amplitudes are induced and demonstrate how they can be tested in experiment and used for quantum devices.

TT 28.2 Wed 15:45 HSZ 105
Broken time-reversal-symmetry in triplet superconductor junctions — ●PHILIP BRYDON¹, CHRISTIAN INIOTAKIS², DIRK MANSKE³, and MANFRED SIGRIST² — ¹Technische Universität Dresden, Dresden, Germany — ²ETH Zürich, Zürich, Switzerland — ³Max-Planck-Institut für Festkörperforschung, Stuttgart, Germany

A rich variety of unconventional Josephson effects have been predicted for junctions combining magnetism and triplet superconductivity (e.g. P. M. R. Brydon *et al.*, Phys. Rev. B **77**, 104504 (2008); P. M. R. Brydon, D. Manske and M. Sigrist, J. Phys. Soc. Japan **77**, 103714 (2008)). Previous works assume, however, that the properties of the barrier material are independent of the two superconductors. We demonstrate that this assumption fails in a scenario where time-reversal symmetry is broken by the misalignment of the **d**-vectors of the triplet superconductors on either side of the junction. This allows the stabilization of a barrier magnetization, creating an exotic Josephson state distinguished by the existence of fractional flux quanta at the junction barrier. There is also a pronounced enhancement of the critical current through the junction at temperatures below the magnetic transition.

TT 28.3 Wed 16:00 HSZ 105
Non-local transport in normal-metal/superconductor hybrid structures: the role of interference and interaction — ●JAKOB BRAUER¹, DETLEF BECKMANN¹, FLORIAN HÜBLER¹, and HILBERT V. LÖHNEYSEN² — ¹Forschungszentrum Karlsruhe, Institut für Nanotechnologie, P.O.-Box 3640, D-76021 Karlsruhe — ²Forschungszentrum Karlsruhe, Institut für Festkörperphysik, P.O.-Box 3640, D-76021 Karlsruhe and Physikalisches Institut, Universität Karlsruhe, D-76128 Karlsruhe, Germany

We present experimental results on non-local conductance in multi-terminal hybrid structures, where two normal metal contacts are attached to a single superconductor. For contacts with an insulating tunnel barrier, and at energies below the energy gap of the superconductor, the non-local conductance is determined by the competition of crossed Andreev reflection (CAR) and elastic cotunneling (EC). The contributions of CAR and EC are expected to cancel each other in the tunneling limit. Recently [Russo *et al.*, Phys. Rev. Lett. **95**, 027002 (2005)], a non-vanishing signal has been observed in such structures,

with an additional energy scale below the gap. So far, quantum interference and Coulomb interaction have been suggested to lift the cancellation of CAR and EC, but no established theory exists for this signal. We observe similar signals in our structures, and demonstrate that the origin is quantum interference.

TT 28.4 Wed 16:15 HSZ 105
Crossed Andreev reflection and dynamical Coulomb blockade — ●ANDREAS BAUMGARTNER, ANDREAS KLEINE, JELENA TRBOVIC, and CHRISTIAN SCHÖNENBERGER — Institute of Physics, University of Basel, Klingelbergstrasse 82, 4056 Basel, Switzerland

A natural source of entangled electrons is the nonlocal process of crossed Andreev reflection (CAR) [1]. In CAR the two electrons of a Cooper pair in a superconductor coherently tunnel into two spatially separated normal metal contacts. This process is expected to produce a negative nonlocal voltage, U_{nl} , in a four terminal device with two normal (injector and detector) and two superconducting contacts. However, recent experiments have shown that elastic cotunneling (EC) and charge imbalance (CI) lead to $U_{nl} > 0$ and can mask CAR [2].

In this contribution we show that U_{nl} can be negative for all subgap biases, which suggests that CAR can dominate all other processes, as required for a solid-state entangler. We fabricated a series of lateral multiterminal Al/Al₂O₃/Pd hybrid structures with contact distances smaller than the superconducting coherence length and with different barrier resistances. We show that for a small window of injector and detector resistances CAR is the dominant nonlocal subgap process, and that for larger resistances the CAR and CI rates are reduced. We tentatively ascribe these systematic changes with barrier resistance to dynamical Coulomb blockade [1].

[1] Recher *et al.*, PRL **91**, 267003 (2003).

[2] Cadden-Zimansky *et al.*, PRL **97**, 237003 (2006), Russo *et al.*, PRL **95**, 027002 (2005), Beckmann *et al.*, PRL **93**, 197003 (2004)

15 min. break

TT 28.5 Wed 16:45 HSZ 105
Hybrid normal-superconducting systems comprising interacting quantum dots — ●MICHELE GOVERNALE¹, MARCO G. PALA², DAVID FUTTERER¹, and JÜRGEN KÖNIG¹ — ¹Theoretische Physik, Universität Duisburg-Essen, D-47048 Duisburg, Germany — ²IMEP-LAHC, INP MINATEC, Centre National de la Recherche Scientifique, F-38016 Grenoble, France

Quantum dots tunnel-coupled to both normal and superconducting leads exhibit a very rich physics due to the presence of superconducting correlations, quantum fluctuations, strong electron-electron interaction, and non-equilibrium. In order to study these systems, we have developed a real-time diagrammatic expansion in the tunnel coupling to the leads [1], which describes both the equilibrium and non-equilibrium superconducting proximity effects in the quantum dot. In the limit of a large superconducting gap, all orders in the tunnel-coupling strength to the superconductors can be included within an exact resummation scheme. Corrections due to finite values of the gap are evaluated within a $1/\Delta$ expansion. This theory is applied to a single-level quantum dot tunnel coupled to two phase-biased superconducting leads and one voltage-biased normal lead. The normal lead is used to drive the dot out of equilibrium. We compute both the Josephson current between the two superconductors and the Andreev current in the normal lead, and analyze their switching on and off as well as transitions between 0- and π -states as a function of gate and bias voltage.

[1] M. Governale, M. G. Pala, and J. König, Phys. Rev. B **77**, 134513 (2008).

TT 28.6 Wed 17:00 HSZ 105